

UM-P-92/69
OZ-92/21
July 1992

Topological phase due to electric dipole moment
and magnetic monopole interaction

Xiao-Gang He and Bruce H.J. McKellar

Research Center for High Energy Physics

School of Physics

University of Melbourne

Parkville, Vic. 3052 Australia

We show that there is an analogous Aharonov-Casher topological effect on a neutral particle with electric dipole moment interacting with a magnetic field produced by magnetic monopoles.

In 1984 Aharonov and Casher(AC) [1] showed that there is a topological effect on a neutral particle with magnetic dipole moment when the particle is moving on a plane under the influence of an electric field produced by a uniformly charged infinitely long filament perpendicular to the plane. When the particle moves around the filament, the wave function will develop a phase equal to $\mu\lambda_e$. Here μ is the magnetic dipole moment of the particle and λ_e is the linear electric charge density of the filament. In the configuration discussed here, the particle is classically force free. This effect is called the AC effect. Several groups have considered this effect [2,3]. A more general derivation and the specific conditions for the AC effect is given in Ref. [2]. The AC effect has also been observed experimentally using neutrons [4].

In this comment we remark that should magnetic monopoles exist, the dual of AC effect would be possible, namely, a neutral particle with an electric dipole moment d will experience a topological effect due to the magnetic field $\mathbf{B}$ produced by magnetic monopoles. The equation of motion for a spin half neutral particle with an electric dipole moment is

$$(\not{\partial} + m + i\frac{d}{2}\sigma_{\mu\nu}\gamma_5 F^{\mu\nu})\psi = 0 . \quad (1)$$

Assuming that there is only a static magnetic field $\mathbf{B}$, this becomes

$$(\not{\partial} + m + id\sigma \cdot \mathbf{B}\gamma_5)\psi = 0 , \quad (2)$$

which can be further rewritten as

$$(\not{\partial} + m - d\gamma \cdot \mathbf{B}\gamma_4)\psi = 0 . \quad (3)$$

This equation has the same form as the equation of motion for a neutral particle with a magnetic dipole moment μ moving in a static electric field $\mathbf{E}$. Indeed changing $-d$ to μ and $\mathbf{B}$ to $\mathbf{E}$ we obtain the equation of motion for the later. We immediately anticipate that equation (2) implies the existence of a topological phase for the particle wavefunction when appropriate conditions are satisfied. One of the conditions for topological AC phase in the wave function is that there are regions where $\nabla \cdot \mathbf{E} \neq 0$. In our case one would require

that there are regions where $\nabla \cdot \mathbf{B}$ is not zero. But in the Maxwellian electrodynamics the magnetic field $\mathbf{B}$ is always divergence-less, and one would not have any topological phase. However, if there are Dirac magnetic monopoles one will have $\nabla \cdot \mathbf{B} = \rho_m$, where ρ_m is magnetic monopole density. Although magnetic monopoles have not been observed it is interesting to speculate along these lines. In the following we will determine the conditions for developing topological phase in the wave function.

Our discussion is modeled on that for the AC effect in Ref. [2]. The basic idea is to find a solution of equation (1) which can be written as

$$\psi = \exp[-ida\mathbf{a}\gamma_4 \int^{\mathbf{r}} \mathbf{\Gamma} \cdot d\mathbf{l}] \psi' , \quad (4)$$

where ψ' is a solution of free Dirac equation of motion, the matrix $\mathbf{a}$ and the field $\mathbf{\Gamma}$ are determined by matching equation(1) with equation(4). One also must make sure that $\nabla \times \mathbf{\Gamma} = 0$ in the region of space where equation (4) holds. We find

$$-d\gamma \cdot \mathbf{B} = i\gamma \cdot \mathbf{\Gamma} \mathbf{a}; \quad \mathbf{a}\gamma_4\gamma_\mu = \gamma_\mu\mathbf{a}\gamma_4. \quad (5)$$

These equations have consistent solutions only in two or less dimensions (spatial). The one dimensional solution is of no interesting to us. So we must consider the motion of the particle move in a plane. Let the plane be the x-y plane. We find $\mathbf{a} = \sigma_{12}$, $\mathbf{\Gamma} = (-B_y, B_x, 0)$ and on this plan $\partial_z B_z = 0$. In the $\nabla \cdot \mathbf{B} = 0$ region, we have

$$\psi = \exp[id\sigma_{12}\gamma_4 \int^{\mathbf{r}} (\mathbf{B} \times \mathbf{z}) \cdot d\mathbf{l}] \psi' , \quad (6)$$

where $\mathbf{z}$ is the unit vector along z-direction. In this region $\nabla \times \mathbf{\Gamma}$ does vanish.

If we let the particle to move a closed path in a plane in the $\nabla \cdot \mathbf{B} = 0$ region which encloses some region where $\nabla \cdot \mathbf{B}$ is not zero, the wave function of the particle will develop a phase

$$\phi = -d\lambda_m , \quad (7)$$

where λ_m is the linear magnetic monopole density. The phase is respect to ψ' . We have shown that there is an analogous AC effect as stated earlier.

There is a 1.5 standard deviation discrepancy between the experimental value and the theoretical prediction for the AC effect. $\phi_{exp}/\phi_{theo} = 1.46 \pm 0.35$ and $\phi_{exp} = 2.19 \pm 0.52 \text{ mrad}$. While we would not wish to encourage the idea that this discrepancy is more than an indication that this difficult experiment should be repeated, we have checked on its implications for the present effect. Taking into account the limit on the electric dipole moment of the neutron ($d < 1.2 \times 10^{-25} \text{ ecm}$ [5]), the linear magnetic monopole density would have to be at least $10^{22}/\text{ecm}$ to bring theory and experiment into agreement. Needless to say, such large magnetic monopole densities are ruled out by other experiments [6].

ACKNOWLEDGMENTS

XGH would like to thank A. Davies, A. Klein and G. Opat for useful discussions. This work is supported in part by the Australian Research Council. We thank the Department of Energy's Institute for Nuclear Theory at the University of Washington for hospitality and the Department of Energy for partial support during the completion of this work.

REFERENCES

- [1] Y. Aharonov and A. Casher, Phys. Rev. Lett. **53**, 319(1984); J. Anandan, Phys. Rev. Lett. **48**, 1660(1982).
- [2] X.-G. He and B. McKellar, Phys. Lett. B**256**, 250(1991).
- [3] A.S. Goldhaber, Phys. Rev. Lett. **62**, **482(1989)**; T. Boyer, Phys. Rev. A**36**, 5083(1987); Y. Aharonov, P. Pearl and L. vaidman, Phys. Rev. A**37**, 4052(1988); C.R. Hagen, Phys. Rev. Lett. **64**, 2347(1990); X.-G. He and B. McKellar, Phys. Lett. B**264**, 129(1991); R. Mignani, J. Phys. A**24**, L421(1991); B. Holstein, Am. J. Phys. **56**, 1080(1991); X.-G. He, in Proceedings of the 26th Rencontre de Moriond: Test of fundamental laws in physics. Les Arcs, Savoie, France. Jan 26 - Feb. 2, 1991.
- [4] A. Cimmino et al., Phys. Rev. Lett. **63**, 380(1989).
- [5] K. Smith et al., Phys. Lett. B**234**, 234(1990).
- [6] NATO ASI Series B: Physics Vol. 111, Monopole' 83, Edited by J. Stone.